

AD 721725

DEPARTMENT
OF
OPERATIONS RESEARCH

GLOBAL STABILITY IN n -PERSON GAMES


by

Louis J. Billera



COLLEGE OF ENGINEERING
CORNELL UNIVERSITY
ITHACA, NEW YORK 14850

Reproduced by
NATIONAL TECHNICAL
INFORMATION SERVICE
SPRINGER-VERLAG, NEW YORK



DEPARTMENT OF OPERATIONS RESEARCH
COLLEGE OF ENGINEERING
CORNELL UNIVERSITY
ITHACA, NEW YORK

TECHNICAL REPORT NO. 127

December 1970

GLOBAL STABILITY IN n -PERSON GAMES

by

Louis J. Billera

This work was supported by the National Science Foundation under grant GK-4795, and partially by the Office of Naval Research under grant N00014-67-A-0077-0014.

Global Stability in n-Person Games

Abstract

A class of bargaining sets, including the bargaining set $M_1^{(i)}$ and the kernel, is treated with regard to studying the tendency to reach stability from unstable points. A known discrete procedure is extended, and these results are applied to derive global stability properties for the solutions of certain differential equations. These differential equations are given in terms of the demand functions which define the bargaining sets, and the set of critical points is precisely the bargaining set in question.

1. INTRODUCTION

Most solution concepts that have been proposed for n -person games are essentially static in nature. That is, once a "stable" point is proposed, there will be no "objection" to that point, and the tendency will be to stay there. However, nothing is said about what is likely to occur at a point which is not stable. There is nothing generally in the definition of stability to guarantee that an unstable situation will tend toward a stable one.

In [14], Stearns showed that for a class of solution concepts related to the bargaining set $M_1^{(i)}$ [10] and the kernel [5], one could define a discrete procedure which, starting with any payoff, converges to a stable point. In this paper, we extend the notion of transfer sequence and, relying heavily on Stearns' methods, prove the necessary convergence results.

We then consider the subclass of Stearns' class of bargaining sets which can be described by continuous demand functions. With each of these bargaining sets we associate a system of differential equations, the critical points of which are precisely the points in the bargaining sets. We prove that for any initial point, there is a solution to this system which approaches a stable point as $t \rightarrow \infty$. In the case of the bargaining set $M_1^{(i)}$ and the kernel, these differential equations have unique solutions for any initial point.

Intuitively, these results have the following interpretation. At each payoff x , there is a number $d_{ij}(x)$ which measures the demand of player i against player j . A point is stable if, for all pairs of players, the demands are zero. The differential equations express the behavior that at any payoff x , all the players are acting to pay off all that they owe, and collect all they are owed. That is,

$$x_i' = \sum_j d_{ij}(x) - \sum_j d_{ji}(x) .$$

It is this behavior that we prove leads to stability.

It is felt that these results are the first steps toward meeting the criticism that n -person game theory ignores the dynamic aspects of the situations it purports to model. It is hoped more results of this type will be forthcoming, in particular in such areas as the core, games without side payments, Nash equilibrium points and games with infinitely many players.

2. GENERAL TRANSFER SEQUENCES

Let n be a positive integer, and let $N = \{1, \dots, n\}$. Let Y be a compact, convex polyhedron in R^n , real Euclidean n -space. We will denote by $R^{n \times n}$ the Euclidean space of points p having n^2 coordinates, denoted p_{ij} for all $i, j \in N$. We will also use the notation $[x]^+ = \max(x, 0)$.

We define the function $\alpha: Y \rightarrow R^{n \times n}$ by the coordinate functions

$$\alpha_{ij}(x) = \begin{cases} \sup\{\epsilon \mid x(\epsilon; i, j) \in Y\} & \text{if } i \neq j \\ 0 & \text{if } i = j, \end{cases}$$

where $x \in Y$, $x(\epsilon; i, j) = (x_1, \dots, x_i + \epsilon, \dots, x_j - \epsilon, \dots, x_n)$. We note here that $\alpha(x)$ is defined and nonnegative (where this means each coordinate is nonnegative) for each $x \in Y$. Since Y is convex, it follows that α is a concave function on Y . It follows from linear programming theory that α is continuous and piecewise linear on Y .

Let v be an n -person game, that is, let $v: 2^N \rightarrow R$ be such that $v(\emptyset) = 0$. Thinking of points in R^{2^N} as being indexed by subsets of N , we define $e: R^n \rightarrow R^{2^N}$ by the coordinate functions

$$e_S(x) = v(S) - x(S),$$

where $x(S)$ is used to denote $\sum_{i \in S} x_i$, $x(\emptyset) = 0$. Now we define $s: R^n \rightarrow R^{n \times n}$ by

$$s_{ij}(x) = \begin{cases} \max(e_T(x) | i \in T, j \notin T) & \text{if } i \neq j \\ 0 & \text{if } i = j. \end{cases}$$

Finally, we define $k: R^n \rightarrow R^{n \times n}$ and $k^*: Y \rightarrow R^{n \times n}$ by

$$k_{ij}(x) = \frac{1}{2}[s_{ij}(x) - s_{ji}(x)]^+ \quad \text{and,}$$

$$k_{ij}^*(x) = \min(k_{ij}(x), a_{ij}(x)) .$$

Note that $k(x) \geq 0$ and $k^*(y) \geq 0$ for $x \in R^n$ and $y \in Y$.

It is easy to see that

$$K(y) = \{x \in Y | k^*(x) = 0\}$$

where $K(Y)$ is the kernel of Y (see [13] for the definitions). In particular, let B be a partition of N and define

$$Y_B = \{x \in R^n \mid x_i \geq v(\{i\}), i \in N; x(S) = v(S), S \in B\}.$$

Then $K(Y_B)$ is the kernel for the coalition structure B studied in [5], [6] and [7]. We remark here that Y_B is a compact, convex polyhedron (which, however, may be empty since we are assuming nothing about the game v).

Let $n: R^{2^n} \rightarrow R^{2^n}$ be the ordering function, that is, the coordinates of $n(z)$ are just the coordinates of z arranged in decreasing order: $n^1(z) \geq n^2(z) \geq \dots \geq n^{2^n}(z)$. We let $\theta: R^n \rightarrow R^{2^n}$ be the composition $n \circ e$, and finally, for each $i = 1, 2, \dots, 2^n$, we let

$$\phi_i: R^n \rightarrow R$$

be such that $\phi_i(x) = \sum_{j=1}^i \theta_j(x)$, for $x \in R^n$. For convenience, we define $\phi_0 \equiv 0$.

Lemma 2.1: Let $x \in R^n$ and let $y = x(\epsilon; i, j)$ where $0 \leq \epsilon \leq k_{ij}(x)$. Then for each $i = 1, \dots, 2^n$, there exists an integer $T < i$ such that

$$\phi_i(y) - \phi_i(x) \leq i(\phi_T(x) - \phi_T(y)).$$

Proof: If $\epsilon = 0$, we may take $T = 0$, so we assume that $\epsilon > 0$. Let

$$A = \{S \subset N \mid e_S(x) \geq s_{ij}(y)\},$$

and let $i^* = |A|$. Suppose $s_{ij}(x) = v_T(x)$ and $s_{ji}(x) = v_R(x)$. From the

definition of k_{ij} , it follows that

$$s_{ij}(x) - \epsilon = s_{ij}(y) \geq s_{ji}(y) = s_{ji}(x) + \epsilon$$

and therefore that $T \in A$, $R \notin A$. Thus for $l < l^*$,

$$\phi_l(y) - \phi_l(x) \leq 0,$$

while

$$\phi_{l^*}(y) - \phi_{l^*}(x) \leq -\epsilon.$$

If $l \leq l^*$, the lemma follows by choosing $\bar{l} = 0$. Since no coordinate of e changes by more than ϵ , we may choose $\bar{l} = l^*$ if $l > l^*$.

Lemma 2.2: Let $x, y^1, \dots, y^m \in R^n$, and suppose that for each $i = 1, \dots, m$ and each $l = 1, \dots, 2^n$, there exists an integer $\bar{l} < l$ such that

$$\phi_l(y^i) - \phi_l(x) \leq l(\phi_{\bar{l}}(x) - \phi_{\bar{l}}(y^i)).$$

Then if $y = \frac{1}{m} \sum_{i=1}^m y^i$, we have for each $l = 1, \dots, 2^n$

$$\phi_l(y) - \phi_l(x) \leq (ml)^l \sum_{0 \leq \bar{l} < l} [\phi_{\bar{l}}(x) - \phi_{\bar{l}}(y)]^+.$$

Proof: We first note that for each ℓ ,

$$\phi_{\ell}(y) \leq \frac{1}{m} \sum_{i=1}^m \phi_{\ell}(y^i) .$$

We may assume without loss of generality that we have equality for each ℓ .

Suppose $\phi_{\ell}(y) - \phi_{\ell}(x) = \omega_0 > 0$. Then there is an index i_0 and an $\ell_1 < \ell$ such that

$$\omega_0 \leq \phi_{\ell}(y^{i_0}) - \phi_{\ell}(x) \leq \ell(\phi_{\ell_1}(x) - \phi_{\ell_1}(y^{i_0})) .$$

Thus

$$\phi_{\ell_1}(x) - \phi_{\ell_1}(y^{i_0}) \geq \frac{\omega_0}{\ell} .$$

Let ω_1 be such that

$$[\phi_{\ell_1}(x) - \phi_{\ell_1}(y)]^+ = \frac{\omega_0}{m\ell} - \frac{\omega_1}{m\ell} .$$

If $\omega_1 > 0$, then there is an i_1 such that

$$\phi_{\ell_1}(y^{i_1}) - \phi_{\ell_1}(x) \geq \frac{\omega_1}{m\ell} .$$

Proceeding inductively, suppose ℓ_{j-1} , i_{j-1} and ω_{j-1} have been defined so that

$$\phi_{\ell_{j-1}}(y^{i_{j-1}}) - \phi_{\ell_{j-1}}(x) \geq \omega_{j-1}/(m\ell)^{j-1} .$$

Then there is an $\ell_j < \ell_{j-1}$ such that

$$\phi_{\ell_j}(x) - \phi_{\ell_j}(x^{i_{j-1}}) \geq \omega_{j-1}/(m\ell)^{j-1}\ell_{j-1} \geq \omega_{j-1}/m^{j-1}\ell^j.$$

Let ω_j be such that

$$[\phi_{\ell_j}(x) - \phi_{\ell_j}(y)]^+ = \frac{\omega_{j-1} - \omega_j}{(m\ell)^j}.$$

If $\omega_j > 0$, there is an i_j such that

$$\phi_{\ell_j}(y^{i_j}) - \phi_{\ell_j}(x) \geq \omega_j/(m\ell)^j.$$

Since $\ell > \ell_1 > \dots > \ell_j \geq 0$, we must eventually reach an integer $p \leq \ell$ such that $\omega_p \leq 0$. Then we have

$$\begin{aligned} (m\ell)^\ell \sum_{0 \leq i < \ell} [\phi_i(x) - \phi_i(y)]^+ &\geq (m\ell)^\ell \sum_{i=1}^p \left(\frac{\omega_{i-1} - \omega_i}{(m\ell)^i} \right) \\ &\geq \sum_{i=1}^p (\omega_{i-1} - \omega_i) = \omega_0 - \omega_p \\ &\geq \omega_0 = \phi_\ell(y) - \phi_\ell(x). \end{aligned}$$

Lemma 2.3: There exists a real number c such that for $x \in \mathbb{R}^n$, if for each $i, j \in \mathbb{N}$, $y^{ij} = x(\epsilon_{ij}; i, j)$ where $0 \leq \epsilon_{ij} \leq k_{ij}(x)$, then if we set $y = \frac{1}{n^2} \sum_{i, j \in \mathbb{N}} y^{ij}$, we have

$$||y-x|| \equiv \sum_{i \in N} |y_i - x_i| \leq c \sum_{l=0}^{2^n} [\phi_l(x) - \phi_l(y)]^+.$$

Proof: Since for $i \in N$,

$$y_i = x_i + \frac{1}{n^2} \sum_{j \in N} (\epsilon_{ij} - \epsilon_{ji}),$$

we have

$$\begin{aligned} ||y-x|| &= \frac{1}{n^2} \sum_{i \in N} \left| \sum_{j \in N} (\epsilon_{ij} - \epsilon_{ji}) \right| \\ &\leq \frac{2}{n^2} \sum_{i,j \in N} \epsilon_{ij} \\ &\leq 2 \max_{i,j \in N} \epsilon_{ij}. \end{aligned}$$

Suppose $\epsilon_{ab} = \max_{i,j \in N} \epsilon_{ij} > 0$. As in the proof of Lemma 2.1, there is an l such that

$$\phi_l(y^{ab}) - \phi_l(x) \leq -\epsilon_{ab}.$$

Let ω_1 be such that

$$[\phi_l(x) - \phi_l(y)]^+ = \frac{\epsilon_{ab} - \omega_1}{n^2}.$$

If $\omega_1 > 0$, then continue as in the proof of Lemma 2.2, to obtain

$$\epsilon_{ab} \leq (n^{2^{2^n}})^{2^n} \sum_{p=0}^{2^n} [\phi_p(x) - \phi_p(y)]^+.$$

Setting $c = 2(n^{2^{2^n}})^{2^n}$, we have the desired result.

Lemma 2.4: Suppose for each $i, j \in N$, we are given ϵ_{ij} where $0 \leq \epsilon_{ij} \leq k_{ij}(x)$ for a fixed $x \in R^n$. Then $\sum_{j \in N} (\epsilon_{ij} - \epsilon_{ji}) = 0$ for each $i \in N$ if and only if $\epsilon_{ij} = 0$ for each $i, j \in N$.

Proof: Sufficiency is obvious. To prove necessity, notice first that if $i, j, l \in N$ then $k_{ij}(x) > 0$ implies $k_{ji}(x) = 0$, and $k_{ij}(x) > 0$ and $k_{jl}(x) > 0$ imply $k_{il}(x) > 0$. The proof of this follows almost exactly as in [5; Lemma 5.1].

Suppose for each $i \in N$, $\sum_{j \in N} (\epsilon_{ij} - \epsilon_{ji}) = 0$, but $\epsilon_{i_1 i_2} > 0$. Thus $k_{i_1 i_2}(x) > 0$. By checking the above sum for $i = i_2$, we find an i_3 so that $\epsilon_{i_2 i_3} > 0$; hence $k_{i_2 i_3}(x) > 0$ which implies $k_{i_1 i_3}(x) > 0$. We may continue this process and obtain i_4, \dots, i_p so that $\epsilon_{i_l i_{l+1}} > 0$ and $k_{i_l i_{l+1}}(x) > 0$, $l = 1, \dots, p-1$. Since the number of indices is finite, we must eventually have some index appear twice. Assume without loss of generality that $i_1 = i_p$. Then $k_{i_1 i_1}(x) > 0$, which is impossible. Therefore $\epsilon_{ij} = 0$ for all $i, j \in N$.

Definition 2.5: Let $x \in R^n$. The x-core, $C(x)$, is given by

$$C(x) = \{y \in R^n \mid y(N) = x(N); \theta_1(y) \leq \theta_1(x)\}.$$

Note that for any x , $C(x)$ is a compact, convex polyhedron and $x \in C(x)$. We remark that if x is a payoff vector, then $C(x)$ coincides with the strong $\theta_1(x)$ -core for $\theta_1(x) \geq 0$, and $C(x)$ is the core when x belongs to the core (i.e., when $\theta_1(x) = 0$). (See [8].) It is easy to see that if $y \in C(x)$ then $C(y) \subset C(x)$.

Definition 2.6: Suppose $x \in R^n$, and for each $i, j \in N$, $0 \leq \epsilon_{ij} \leq \frac{1}{2} k_{ij}(x)$. Let $y \in R^n$ be defined by

$$y_i = x_i + \sum_{j \in N} (\epsilon_{ij} - \epsilon_{ji}) \quad \text{for } i \in N.$$

Then we say y results from x by a k-bounded transfer.

We observe here that if y results from x by a k -bounded transfer, then if we set $\delta_{ij} = n^2 \epsilon_{ij}$, we have $y = \frac{1}{n^2} \sum_{i,j \in N} x(\delta_{ij}; i, j)$. Thus, y satisfies the hypotheses of Lemmas 2.2 and 2.3.

Lemma 2.7: If y results from x by a k-bounded transfer, then $y \in C(x)$.

Proof: It follows from the definition that $y(N) = x(N)$. Since $\phi_1 = \theta_1$, the result follows from Lemma 2.2.

Defining k^* -bounded transfers in the obvious way, we can observe that if $x \in Y$ and y results from x by a k^* -bounded transfer, then $y \in C(x) \cap Y$.

We say a sequence $\{x^i\}_{i=0}^{\infty}$ of points in R^n is a transfer sequence if for each i , x^{i+1} results from x^i by a k -bounded transfer. By applying Lemma 2.7, we can obtain the following result.

Lemma 2.8: If $\{x^i\}_{i=0}^{\infty}$ is a transfer sequence, then for each i , $x^i \in C(x^0)$. Thus every transfer sequence is bounded. Further, if $x^0 \in Y$ and all the transfers are k^* -bounded, then $x^i \in C(x^0) \cap Y$ for each i .

We are able to say more. As we will next prove, every transfer sequence is convergent.

Theorem 2.9: Given $x^0 \in R^n$, there exists a real number $\beta(x^0)$, such that for all transfer sequences $\{x^i\}_{i=0}^{\infty}$ starting at x^0 ,

$$\sum_{i=0}^{\infty} \|x^{i+1} - x^i\| \leq \beta(x^0). \quad (2.9.1)$$

Further, given a compact, convex set Y , there exists a real number $\beta(Y)$, independent of the game v , such that for any $x^0 \in Y$ and any sequence $\{x^i\}_{i=0}^{\infty}$ consisting only of k^* -bounded transfers, (2.9.1) is true with $\beta(Y)$ in place of $\beta(x^0)$.

Proof: For each i and integer l , $0 \leq l \leq 2^n$, set

$$\delta_l^+(i) = [\phi_l(x^{i+1}) - \phi_l(x^i)]^+, \quad \text{and}$$

$$\delta_l^-(i) = [\phi_l(x^i) - \phi_l(x^{i+1})]^+.$$

In addition let $I_l^r = \sum_{i=0}^r \delta_l^+(i)$ and $D_l^r = \sum_{i=0}^r \delta_l^-(i)$. Then by Lemma 2.3,

$$||x^{i+1} - x^i|| \leq c \sum_{l=0}^{2^n} \delta_l^-(i),$$

and thus

$$\sum_{i=0}^r ||x^{i+1} - x^i|| \leq c \sum_{l=0}^{2^n} D_l^r.$$

We will show the existence of a $\beta(x^0)$ (or $\beta(Y)$, in the second case) such that for all $r \geq 0$

$$c \sum_{l=0}^{2^n} D_l^r \leq \beta(x^0) \quad (\text{or } \beta(Y)).$$

By Lemma 2.2, we have for each l ,

$$\delta_l^+(i) \leq (n^2 l)^l \sum_{0 \leq j < l} \delta_j^-(i).$$

Therefore, for all $r \geq 0$, by summing from 0 to r , we get

$$I_l^r \leq (n^2 l)^l \sum_{0 \leq j < l} D_j^r. \quad (2.9.2)$$

Now $\delta_l^-(i) - \delta_l^+(i) = \phi_l(x^i) - \phi_l(x^{i+1})$, so

$$D_l^r - I_l^r = \phi_l(x^0) - \phi_l(x^{r+1}). \quad (2.9.3)$$

By Lemma 2.8, both $x^0, x^{r+1} \in C(x^0)$ (or $C(x^0) \cap Y$, for the second case).

By the compactness of $C(x^0)$ (or Y), there exists a $\rho(x^0)$ ($\rho(Y)$) such that

$$\max_{S \subseteq N} |x(S) - y(S)| \leq \rho(x^0) \quad (\rho(Y))$$

for all $x, y \in C(x^0)$ ($x, y \in Y$). Thus for $S \subseteq N$, $e_S(y) - e_S(x) \leq \rho(x^0)$, and thus $\phi_\ell(y) - \phi_\ell(x) \leq \ell \rho(x^0)$ for $x, y \in C(x^0)$ (or $\phi_\ell(y) - \phi_\ell(x) \leq \ell \rho(Y)$ for $x, y \in Y$). Applying this to (2.9.3), we get for each ℓ

$$D_\ell^r \leq I_\ell^r + \ell \rho(x^0) \quad (\ell \rho(Y)) . \quad (2.9.4)$$

Note here that $\rho(x^0)$ depends on $C(x^0)$, therefore on x^0 and v .

On the other hand, $\rho(Y)$ depends only on Y .

To derive the bounds $\beta(x^0)$ and $\beta(Y)$, notice first that for all $r \geq 0$, $I_0^r = D_0^r = 0$. Now suppose that I_j^r and D_j^r have been given bounds independent of r for $j = 0, \dots, \ell-1$. First (2.9.2) provides a bound for I_ℓ^r , and then (2.9.4) provides a bound for D_ℓ^r . Both these bounds are independent of r . The theorem follows.

3. DEMAND FUNCTIONS AND BARGAINING SETS

By a demand function, we will mean a continuous function $d: R^n \rightarrow R^{n \times n}$ which satisfies

$$0 \leq d(x) \leq k(x)$$

for all $x \in R^n$.

For a given demand function d , we define a function $D: R^n \rightarrow R^n$ by $D_i = \sum_{j \in N} (d_{ij} - d_{ji})$. It follows from Lemma 2.4 that $D(x) = 0$ if and only if $d_{ij}(x) = 0$ for all $i, j \in N$. We now can define the bargaining set associated with the demand function d as

$$M(d) = \{x \in R^n \mid D(x) = 0\}.$$

This agrees with the definition given by Stearns in [14].

Suppose, in addition, that $d(y) \leq k^*(y)$ for $y \in Y$. Then we say d is k^* bounded and we define

$$M_Y(d) = M(d) \cap Y.$$

The Brouwer fixed point theorem shows that $M_Y(d) \neq \emptyset$ since the map $f: Y \rightarrow Y$ given by $f(x) = x + \frac{1}{2n} D(x)$ must have a fixed point. The non-emptiness of $M(d)$ for general d will follow from the next result.

Theorem 3.1: Let $x^0 \in R^n$ and suppose a sequence $(x^i)_{i=0}^\infty$ is defined by $x^{i+1} = x^i + hD(x^i)$ where $0 < h \leq \frac{1}{2n}$. Then (x^i) converges to a point $x \in M(d)$.

Further, if $x^0 \in Y$ and d is k^* bounded, then (x^i) converges to $x \in M_Y(d)$.

Proof: In either case, the sequence is a transfer sequence, and therefore Theorem 2.9 applies to give

$$\sum_{i=0}^{\infty} \|x^{i+1} - x^i\| < \infty.$$

This implies the sequence is Cauchy, since for $m > k$,

$$||x^m - x^k|| \leq \sum_{i=k}^{m-1} ||x^{i+1} - x^i|| \leq \sum_{i=k}^{\infty} ||x^{i+1} - x^i||,$$

which can be made small for large k . Thus there exists $x \in R^n$ s.t. $x^i \rightarrow x$. Since D is continuous, and

$$||D(x^i)|| = \frac{1}{h} ||x^{i+1} - x^i|| \rightarrow 0,$$

it follows that $D(x) = 0$ and $x \in M(d)$ in the first case.

In the second case, Lemma 2.8 implies that $x \in Y$ and by the above argument, $D(x) = 0$.

Corollary 3.2: $M(d) \neq \emptyset$, in fact, for any $x^0 \in R^n$, there exists an $x \in M(d) \cap C(x^0)$. If d is k^* bounded, then for any $x^0 \in Y$, there exists an $x \in M_Y(d) \cap C(x^0)$.

We remark here that in either of the cases treated in 3.1 we have

$$\sum_{i=0}^{\infty} ||x^{i+1} - x^i|| < \beta,$$

where β will depend either on x^0 or on Y , but β is independent of h .

4. GLOBAL STABILITY

Let d be a demand function, and consider the system of differential equations

$$x' = D(x) . \quad (4.1)$$

Since D is continuous, there exist solutions to 4.1 through any initial point $x^0 \in \mathbb{R}^n$. From [9, Theorem 1.31, p. 9], it follows that the solutions of 4.1 are defined for all $t \in \mathbb{R}$, since each D_i is nonnegative and bounded above by the piecewise linear function $\sum_{j \in N} (k_{ij} + k_{ji})$.

We will be concerned here only with solutions defined on $\mathbb{R}_+ = [0, \infty)$ and starting at some fixed initial point, $x^0 \in \mathbb{R}^n$. That is, we will consider functions $\phi: \mathbb{R}_+ \rightarrow \mathbb{R}^n$ such that $\phi(0) = x^0$ and

$$\frac{d\phi_i(t)}{dt} = D_i(\phi(t)) ,$$

for all $t \in \mathbb{R}_+$.

Theorem 4.1: For any $x^0 \in \mathbb{R}^n$, there exists a solution ϕ to $x' = D(x)$ such that $\phi(0) = x^0$ and $\phi(t)$ converges to a point $x \in M(d)$ as $t \rightarrow \infty$. If, in addition, $x^0 \in Y$ and d is k^* bounded, then $\phi(t) \in Y$ for all $t \in \mathbb{R}_+$ and $x \in M_Y(d)$.

Proof: Let $x^0 \in \mathbb{R}^n$, and choose a real number M so that $\|D(x)\| \leq M$ for $x \in C(x^0)$.

Let $0 < h \leq h_0 = \frac{1}{n}$, and define the sequence $\{x^i\}$ by $x^{i+1} = x^i + hD(x^i)$. Notice that $\{x^i\} \subset C(x^0)$, and if $x^0 \in Y$ and d is

k^* bounded, then $\{x^i\} \subset C(x^0) \cap Y$, by Lemma 2.8. Now define

$$\phi^h: R_+ \rightarrow C(x^0)$$

by $\phi^h(0) = x^0$, and

$$\phi^h(t) = x^i + D(x^i)(t - ih)$$

where i is chosen so that $ih < t \leq (i+1)h$. That ϕ^h maps into $C(x^0)$ follows from the above comments and the convexity of $C(x^0)$. Again, if $x^0 \in Y$ and d is k^* bounded, then ϕ^h maps into $C(x^0) \cap Y$.

Let $\langle h \rangle = \{ih | i = 0, 1, 2, \dots\}$. On $R_+ - \langle h \rangle$ we have that $\left(\phi^h\right)'$ exists.* In fact, for $ih < t < (i+1)h$, $\left(\phi^h\right)'(t) = D(x^i)$.

Let $\epsilon > 0$, and choose $0 < h_\epsilon \leq h_0$ so that for $x, y \in C(x^0)$, $\|y - x\| < Mh_\epsilon$ implies $\|D(y) - D(x)\| < \epsilon$. Let $0 < h < h_\epsilon$, $t \in R_+ - \langle h \rangle$, and suppose $ih < t < (i+1)h$. Then since

$$\|\phi^h(t) - x^i\| = \|D(x^i)\| |t - ih| < Mh_\epsilon,$$

we have

$$\|\left(\phi^h\right)'(t) - D(\phi^h(t))\| = \|D(x^i) - D(\phi^h(t))\| < \epsilon.$$

Clearly $\|\phi^h(t) - \phi^h(t')\| \leq M|t - t'|$ for all $0 < h \leq h_0$. Thus the family of functions $\{\phi^h | 0 < h \leq h_0\}$ is uniformly bounded and equicontinuous. Let $\{h_j\}_{j=1}^\infty$ be such that $h_j \leq h_0$ for each j and $h_j \rightarrow 0$

* $\left(\phi^h\right)' = \frac{d\phi^h}{dt}$.

as $j \rightarrow \infty$. Denote ϕ^{h_j} by ϕ^j . By the Ascoli theorem, there is a subsequence of the ϕ^j (we may assume now without loss of generality that it is the entire sequence ϕ^j) and a continuous function

$$\phi: R_+ \rightarrow C(x^0)$$

such that $\phi^j \rightarrow \phi$ pointwise on R_+ as $j \rightarrow \infty$ and the convergence is uniform on compact subsets of R_+ . We show first that ϕ is a solution to $x' = D(x)$. Clearly $\phi(0) = x^0$. Note that $||\phi(t) - \phi(t')|| \leq M|t-t'|$.

Let $\Delta^j(t) = \left\{ \phi^j \right\}'(t) - D \circ \phi^j(t)$ for $t \in R_+$ such that $\left\{ \phi^j \right\}'(t)$ exists. Otherwise let $\Delta^j(t) = 0$. Thus, $D \circ \phi^j(t) + \Delta^j(t) = \left\{ \phi^j \right\}'(t)$ whenever $\left\{ \phi^j \right\}'(t)$ exists, that is, except for a countable subset of R_+ . Since ϕ^j is absolutely continuous, it follows that

$$\phi^j(t) = x^0 + \int_0^t [D \circ \phi^j(\tau) + \Delta^j(\tau)] d\tau.$$

Since $\phi^j \rightarrow \phi$ uniformly on $[0, t]$ and D is uniformly continuous on $C(x^0)$, we have that $D \circ \phi^j \rightarrow D \circ \phi$ uniformly on $[0, t]$. From the discussion above, it follows that $\Delta^j(t) \rightarrow 0$ uniformly on R_+ . Thus

$$\phi(t) = x^0 + \int_0^t D \circ \phi(\tau) d\tau,$$

and finally

$$\phi'(t) = D(\phi(t)).$$

We show now that $\phi(t)$ converges to a point $x \in M(d)$ as $t \rightarrow \infty$.

It follows from the remark following Corollary 3.2 that there is a number β (independent of h) such that the total variation of the function ϕ^h , $0 < h \leq h_0$, on the interval $[0, T]$ is less than β for all $T \in \mathbb{R}_+$. Since $\phi^j \rightarrow \phi$ it follows that the total variation of ϕ on $[0, T]$ is also less than β for all T . Therefore $\phi(t)$ converges to a unique limit point $x \in C(x^0)$ as $t \rightarrow \infty$.

To see that $x \in M(d)$, note that $D\phi(t) \rightarrow D(x)$ as $t \rightarrow \infty$. For each integer $\ell \geq 0$, and for each $i = 1, \dots, n$, there is a $\tau_i^\ell \in \mathbb{R}_+$, such that $\ell \leq \tau_i^\ell \leq \ell+1$ and

$$\begin{aligned} \phi_i(\ell+1) - \phi_i(\ell) &= \phi_i'(\tau_i^\ell) \\ &= D_i \circ \phi(\tau_i^\ell). \end{aligned}$$

Since for each i , $\tau_i^\ell \rightarrow \infty$ as $\ell \rightarrow \infty$, we have

$$D_i \circ \phi(\tau_i^\ell) \rightarrow D_i(x) \text{ as } \ell \rightarrow \infty.$$

But $\phi_i(\ell+1) - \phi_i(\ell) \rightarrow 0$ as $\ell \rightarrow \infty$ for each i , thus $D(x) = 0$ and $x \in M(d)$.

If $x^0 \in Y$ and d is k^* bounded, we have $\phi(\mathbb{R}_+) \subset C(x^0) \cap Y$ since this holds for each ϕ^h . It is clear that x must be in $M_Y(d)$ in this case. This completes the proof.

We remark that in the case where $x^0 \in Y$ and the demand functions are k^* bounded, it is enough to consider them as if they were defined only on Y . This causes no loss of generality since any function

$d^*: Y \rightarrow R^{n \times n}$ satisfying $d^* \leq k^*$ can be extended to a k^* bounded demand function $d: R^n \rightarrow R^{n \times n}$ such that $d|_Y = d^*$.

5. SOME SPECIAL CASES

If the demand function d has the property that the system $x' = D(x)$ has a unique solution through each initial point x^0 , then Theorem 4.1 becomes particularly useful. A simple condition which guarantees uniqueness is that each of the functions d_{ij} is piecewise linear with a finite number of different linear "pieces". Indeed, this condition implies that $D(x)$ is uniformly Lipschitz and hence that solutions of $x' = D(x)$ are unique (see [4]).

It is clear that the function k is piecewise linear on R^n , and since Y is a polyhedron, k^* is piecewise linear on Y . Letting $Y = Y_B$ for some partition B of N (and assuming $Y \neq \emptyset$), we have $K(Y) = N_Y(k^*)$ which is the kernel for the coalition structure B .

Following Stearns [14], we can define a k^* bounded demand function m on Y_B so that $N_Y(m)$ is the bargaining set $N_1^{(i)}$ for the coalition structure B (see [10]). It can be shown that m is also piecewise linear. The continuity of m follows from [1; Lemma 2.11]. Taking these facts together with Theorem 4.1, we can state the following.

Theorem 5.1: The differential equations which describe (in the sense of Section 4) the kernel K and the bargaining set $N_1^{(i)}$ for any partition B have unique solutions through any point $x^0 \in Y_B$. Further, these solutions approach K and $N_1^{(i)}$ as $t \rightarrow \infty$.

In the case where v is monotonic in the zero normalization (see [8]) we can simplify the differential equations which describe the kernel K for the grand coalition (i.e. $B = \{N\}$).

Proposition 5.2: Let $Y = Y_{\{N\}}$. Then if v is monotonic in the zero normalization, we have

$$K = M_Y(k^*) = M(k) \cap \{x \in R^n \mid x(N) = v(N)\}.$$

Proof: That $K = M_Y(k^*)$ is clear. We prove the second equality. The inclusion

$$M_Y(k^*) \subset M(k) \cap \{x \in R^n \mid x(N) = v(N)\}$$

is Lemma 3.4 of [8]. To show the opposite inclusion,* suppose $x(N) = v(N)$ and $s_{ij}(x) = s_{ji}(x)$ for all $i, j \in N$, but $x_k < v(\{k\})$ for some k . Let \mathcal{D} be the set of coalitions S for which $e_S(x)$ is maximum. Since $e_{\{k\}}(x) > 0$, $N \notin \mathcal{D}$. Further, $S \in \mathcal{D}$ implies $k \in S$ since $T \in \mathcal{D}$, $k \notin T$ implies $e_R(x) > e_T(x)$, where $R = T \cup \{k\}$, by monotonicity in the zero normalization and the assumption that $x_k < v(\{k\})$. Therefore, there exists an $\ell \in N$ s.t. $s_{k\ell}(x) > s_{\ell k}(x)$, which is impossible. Thus, $x_k \geq v(\{k\})$.

Theorem 5.3: Let v be monotonic in the zero normalization and let K be the kernel of v for the grand coalition structure. Suppose $x^0 \in R^n$ and $x^0(N) = v(N)$. Then the unique solution ϕ to $x' = K(x)$ such that $\phi(0) = x^0$ has the property that

*The author is indebted to Elon Kohlberg for this proof.

$$\lim_{t \rightarrow \infty} \phi(t) \in K.$$

Proof: This is immediate from Theorem 4.1 and Proposition 5.1.

6. COMMENTS

One problem left by this study is to obtain a direct proof of Theorem 4.1, that is, one which does not require going through discrete transfer methods (at least in the case where the differential equations have unique solutions.) It would seem from the methods of this study that one should be able to obtain stability theorems for systems of differential equations by use of lexicographically decreasing vector "Lyapunov functions." The function θ seems to serve this purpose here. However, the fact that the lexicographic ordering is not continuous keeps the standard proofs from working in this case.

It would be of great interest in Mathematical Economics if global stability results could be obtained for the core. Such results would say something about the tendency of an economy to reach equilibrium. A global stability result for the nucleolus (see [13]) would essentially be a result for the core since the nucleolus is in the core whenever the core is non-empty. We feel it is likely that the results here can be extended in some way to obtain convergence to the nucleolus.

It would also be interesting to extend the notion of a "behavioral vector field" to games without side payments and games with infinitely many players. In the case of infinitely many players, it has so far been difficult even to extend the notion of bargaining set, much less to worry about existence. Perhaps the differential equations approach will prove more

fruitful. Results on games without side payments ([11], [2] and [1]) seem to indicate that static stability may not always be possible, and that we may have to settle for some sort of cyclic behavior. Perhaps this can be formalized in terms of periodic solutions to related differential equations.

Finally, one can describe the Nash equilibrium point solutions to a multiperson cooperative game by demand functions, and, therefore, by a system of differential equations. These equations resemble (but are essentially different than) the differential equations studied by Brown and von Neumann [3] for solving zero-sum two-person games. The notion of demand functions for equilibrium points is motivated by Peleg's treatment of equilibrium points as solutions to open acyclic relations in [12].

Roughly, let S_k be the (finite) set of pure strategies for player k , $k = 1, \dots, m$. Let $n = \sum_{k=1}^m |S_k|$, and let

$$Y = \{x \in \mathbb{R}^n \mid x_i \geq 0; \sum_{i \in S_k} x_i = 1, \quad k = 1, \dots, m\}$$

be the set of vectors of mixed strategies for the m players. For each $k = 1, \dots, m$, let

$$H_k: Y \rightarrow \mathbb{R}$$

be the payoff function for player k and let $H_k(x|i)$ denote the payoff to player k when every other player plays his mixed strategy as given in x , but player k uses his pure strategy i . Let $i, j \in N = \bigcup_{k=1}^m S_k$. We

define the demand function

$$d: Y \rightarrow R^{n \times n}$$

by $d_{ij}(x) = x_j [H_k(x|i) - H_k(x|j)]^+$ when $i, j \in S_k$ for some k . Otherwise $d_{ij}(x) = 0$. It is clear that x is an equilibrium point if and only if $d_{ij}(x) = 0$ for all $i, j \in N$. It is not known whether Theorem 4.1 is true for the system $x' = D(x)$ defined by the function d .

The author is grateful to M. Balch, M. Brown, C. Dafermos and M. Maschler for many helpful conversations on this work.

REFERENCES

- [1] N. Asscher, A bargaining set for cooperative games without side payments, Research Prog. in Game Theory and Math. Eco., Dept. of Math., The Hebrew University of Jerusalem, RM No. 58, April, 1970.
- [2] L. J. Billera, Existence of general bargaining sets for cooperative games without side payments, Bull. Amer. Math. Soc. 76(1970), pp. 375-379.
- [3] G. W. Brown and J. von Neumann, Solutions of games by differential equations, Contributions to the Theory of Games, Vol. I, A. W. Kuhn and A. W. Tucker, eds., Annals of Mathematics Studies, No. 24 (1950), pp. 73-79, Princeton University Press, Princeton, New Jersey.
- [4] E. A. Coddington and N. Levinson, Theory of Ordinary Differential Equations, McGraw-Hill, New York, 1955.
- [5] M. Davis and M. Maschler, The kernel of a cooperative game, Naval Res. Logist. Quart. 12 (1965), pp. 223-259.
- [6] M. Maschler and B. Peleg, A characterization, existence proof and dimension bounds for the kernel of a game, Pacific J. Math., 18 (1966), pp. 289-328.
- [7] _____, The structure of the kernel of a cooperative game, SIAM J. Appl. Math. 15 (1967), pp. 569-604.
- [8] M. Maschler, B. Peleg and L. S. Shapley, The kernel and the nucleolus of a cooperative game as locuses in the strong ϵ -core, Research Prog. in Game Theory and Math. Eco., Dept. of Math., The Hebrew University of Jerusalem, RM No. 60, May, 1970.
- [9] V. V. Nemytskii and V. V. Stepanov, Qualitative Theory of Differential Equations, Princeton University Press, Princeton, New Jersey, 1960.
- [10] B. Peleg, Existence theorem for the bargaining set $M_1^{(i)}$, in Essays in Mathematical Economics, M. Shubik (Editor), Princeton Univ. Press, Princeton, N.J., 1967, cf. Bull. Amer. Math. Soc. 69 (1963), 109-110.
- [11] _____, Bargaining sets of cooperative games without side payments, Israel J. Math. 1 (1967), 197-200.
- [12] _____, Equilibrium points for open acyclic relations, Canad. J. Math. 19 (1967), 366-369.
- [13] D. Schmeidler, The nucleolus of a characteristic function game, SIAM J. Appl. Math. 17 (1969), pp. 1163-1170.
- [14] R. E. Stearns, Convergent transfer schemes for n-person games, Trans. Amer. Math. Soc. 134 (1968), pp. 449-459.

[illegible]

NOTE: The following information must be entered when the overall report is classified:

28. REPORT SECURITY CLASSIFICATION:

2b GROUP

4. Type of report and inclusive dates

* Add CODE suffix first name, middle initial, last name

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60 61 62 63 64 65 66 67 68 69 70 71 72 73 74 75 76 77 78 79 80 81 82 83 84 85 86 87 88 89 90 91 92 93 94 95 96 97 98 99 100 101 102 103 104 105 106 107 108 109 110 111 112 113 114 115 116 117 118 119 120 121 122 123 124 125 126 127 128 129 130 131 132 133 134 135 136 137 138 139 140 141 142 143 144 145 146 147 148 149 150 151 152 153 154 155 156 157 158 159 160 161 162 163 164 165 166 167 168 169 170 171 172 173 174 175 176 177 178 179 180 181 182 183 184 185 186 187 188 189 190 191 192 193 194 195 196 197 198 199 200 201 202 203 204 205 206 207 208 209 210 211 212 213 214 215 216 217 218 219 220 221 222 223 224 225 226 227 228 229 230 231 232 233 234 235 236 237 238 239 240 241 242 243 244 245 246 247 248 249 250 251 252 253 254 255 256 257 258 259 260 261 262 263 264 265 266 267 268 269 270 271 272 273 274 275 276 277 278 279 280 281 282 283 284 285 286 287 288 289 290 291 292 293 294 295 296 297 298 299 300 301 302 303 304 305 306 307 308 309 310 311 312 313 314 315 316 317 318 319 320 321 322 323 324 325 326 327 328 329 330 331 332 333 334 335 336 337 338 339 340 341 342 343 344 345 346 347 348 349 350 351 352 353 354 355 356 357 358 359 360 361 362 363 364 365 366 367 368 369 370 371 372 373 374 375 376 377 378 379 380 381 382 383 384 385 386 387 388 389 390 391 392 393 394 395 396 397 398 399 400 401 402 403 404 405 406 407 408 409 410 411 412 413 414 415 416 417 418 419 420 421 422 423 424 425 426 427 428 429 430 431 432 433 434 435 436 437 438 439 440 441 442 443 444 445 446 447 448 449 450 451 452 453 454 455 456 457 458 459 460 461 462 463 464 465 466 467 468 469 470 471 472 473 474 475 476 477 478 479 480 481 482 483 484 485 486 487 488 489 490 491 492 493 494 495 496 497 498 499 500 501 502 503 504 505 506 507 508 509 510 511 512 513 514 515 516 517 518 519 520 521 522 523 524 525 526 527 528 529 530 531 532 533 534 535 536 537 538 539 540 541 542 543 544 545 546 547 548 549 550 551 552 553 554 555 556 557 558 559 560 561 562 563 564 565 566 567 568 569 570 571 572 573 574 575 576 577 578 579 580 581 582 583 584 585 586 587 588 589 590 591 592 593 594 595 596 597 598 599 600 601 602 603 604 605 606 607 608 609 610 611 612 613 614 615 616 617 618 619 620 621 622 623 624 625 626 627 628 629 630 631 632 633 634 635 636 637 638 639 640 641 642 643 644 645 646 647 648 649 650 651 652 653 654 655 656 657 658 659 660 661 662 663 664 665 666 667 668 669 670 671 672 673 674 675 676 677 678 679 680 681 682 683 684 685 686 687 688 689 690 691 692 693 694 695 696 697 698 699 700 701 702 703 704 705 706 707 708 709 710 711 712 713 714 715 716 717 718 719 720 721 722 723 724 725 726 727 728 729 730 731 732 733 734 735 736 737 738 739 740 741 742 743 744 745 746 747 748 749 750 751 752 753 754 755 756 757 758 759 760 761 762 763 764 765 766 767 768 769 770 771 772 773 774 775 776 777 778 779 780 781 782 783 784 785 786 787 788 789 790 791 792 793 794 795 796 797 798 799 800 801 802 803 804 805 806 807 808 809 810 811 812 813 814 815 816 817 818 819 820 821 822 823 824 825 826 827 828 829 830 831 832 833 834 835 836 837 838 839 840 841 842 843 844 845 846 847 848 849 850 851 852 853 854 855 856 857 858 859 860 861 862 863 864 865 866 867 868 869 870 871 872 873 874 875 876 877 878 879 880 881 882 883 884 885 886 887 888 889 890 891 892 893 894 895 896 897 898 899 900 901 902 903 904 905 906 907 908 909 910 911 912 913 914 915 916 917 918 919 920 921 922 923 924 925 926 927 928 929 930 931 932 933 934 935 936 937 938 939 940 941 942 943 944 945 946 947 948 949 950 951 952 953 954 955 956 957 958 959 960 961 962 963 964 965 966 967 968 969 970 971 972 973 974 975 976 977 978 979 980 981 982 983 984 985 986 987 988 989 990 991 992 993 994 995 996 997 998 999 1000 1001 1002 1003 1004 1005 1006 1007 1008 1009 1010 1011 1012 1013 1014 1015 1016 1017 1018 1019 1020 1021 1022 1023 1024 1025 1026 1027 1028 1029 1030 1031 1032 1033 1034 1035 1036 1037 1038 1039 1040 1

7a. TOTAL NO OF PAGES

| 70. NO OF REFS |
|----------------|
| 1 |
| 2 |
| 3 |
| 4 |
| 5 |
| 6 |
| 7 |
| 8 |
| 9 |
| 10 |
| 11 |
| 12 |
| 13 |
| 14 |
| 15 |
| 16 |
| 17 |
| 18 |
| 19 |
| 20 |
| 21 |
| 22 |
| 23 |
| 24 |
| 25 |
| 26 |
| 27 |
| 28 |
| 29 |
| 30 |
| 31 |
| 32 |
| 33 |
| 34 |
| 35 |
| 36 |
| 37 |
| 38 |
| 39 |
| 40 |
| 41 |
| 42 |
| 43 |
| 44 |
| 45 |
| 46 |
| 47 |
| 48 |
| 49 |
| 50 |
| 51 |
| 52 |
| 53 |
| 54 |
| 55 |
| 56 |
| 57 |
| 58 |
| 59 |
| 60 |
| 61 |
| 62 |
| 63 |
| 64 |
| 65 |
| 66 |
| 67 |
| 68 |
| 69 |
| 70 |
| 71 |
| 72 |
| 73 |
| 74 |
| 75 |
| 76 |
| 77 |
| 78 |
| 79 |
| 80 |
| 81 |
| 82 |
| 83 |
| 84 |
| 85 |
| 86 |
| 87 |
| 88 |
| 89 |
| 90 |
| 91 |
| 92 |
| 93 |
| 94 |
| 95 |
| 96 |
| 97 |
| 98 |
| 99 |
| 100 |

$$u, f \quad p_d \quad \text{to} \quad \Delta \quad \text{to} \quad \text{to} \quad \text{to} \quad \Delta \quad p_d \quad \text{to} \quad \Delta$$

4a. ORIGINATOR'S REPORT NUMBER(S)

1. 6 4 2 0 8 6 4 2 0

IF OTHER REPORT NO(S) (Any other numbers that may be assigned this report)

1. INSTRUCTIONS STATEMENT:

11. 12. 13. 14. 15. 16. 17. 18. 19. 20. 21. 22. 23. 24. 25. 26. 27. 28. 29. 30. 31. 32. 33. 34. 35. 36. 37. 38. 39. 40. 41. 42. 43. 44. 45. 46. 47. 48. 49. 50. 51. 52. 53. 54. 55. 56. 57. 58. 59. 60. 61. 62. 63. 64. 65. 66. 67. 68. 69. 70. 71. 72. 73. 74. 75. 76. 77. 78. 79. 80. 81. 82. 83. 84. 85. 86. 87. 88. 89. 90. 91. 92. 93. 94. 95. 96. 97. 98. 99. 100. 101. 102. 103. 104. 105. 106. 107. 108. 109. 110. 111. 112. 113. 114. 115. 116. 117. 118. 119. 120. 121. 122. 123. 124. 125. 126. 127. 128. 129. 130. 131. 132. 133. 134. 135. 136. 137. 138. 139. 140. 141. 142. 143. 144. 145. 146. 147. 148. 149. 150. 151. 152. 153. 154. 155. 156. 157. 158. 159. 160. 161. 162. 163. 164. 165. 166. 167. 168. 169. 170. 171. 172. 173. 174. 175. 176. 177. 178. 179. 180. 181. 182. 183. 184. 185. 186. 187. 188. 189. 190. 191. 192. 193. 194. 195. 196. 197. 198. 199. 200. 201. 202. 203. 204. 205. 206. 207. 208. 209. 210. 211. 212. 213. 214. 215. 216. 217. 218. 219. 220. 221. 222. 223. 224. 225. 226. 227. 228. 229. 230. 231. 232. 233. 234. 235. 236. 237. 238. 239. 240. 241. 242. 243. 244. 245. 246. 247. 248. 249. 250. 251. 252. 253. 254. 255. 256. 257. 258. 259. 260. 261. 262. 263. 264. 265. 266. 267. 268. 269. 270. 271. 272. 273. 274. 275. 276. 277. 278. 279. 280. 281. 282. 283. 284. 285. 286. 287. 288. 289. 290. 291. 292. 293. 294. 295. 296. 297. 298. 299. 300. 301. 302. 303. 304. 305. 306. 307. 308. 309. 310. 311. 312. 313. 314. 315. 316. 317. 318. 319. 320. 321. 322. 323. 324. 325. 326. 327. 328. 329. 330. 331. 332. 333. 334. 335. 336. 337. 338. 339. 340. 341. 342. 343. 344. 345. 346. 347. 348. 349. 350. 351. 352. 353. 354. 355. 356. 357. 358. 359. 360. 361. 362. 363. 364. 365. 366. 367. 368. 369. 370. 371. 372. 373. 374. 375. 376. 377. 378. 379. 380. 381. 382. 383. 384. 385. 386. 387. 388. 389. 390. 391. 392. 393. 394. 395. 396. 397. 398. 399. 400. 401. 402. 403. 404. 405. 406. 407. 408. 409. 410. 411. 412. 413. 414. 415. 416. 417. 418. 419. 420. 421. 422. 423. 424. 425. 426. 427. 428. 429. 430. 431. 432. 433. 434. 435. 436. 437. 438. 439. 440. 441. 442. 443. 444. 445. 446. 447. 448. 449. 450. 451. 452. 453. 454. 455. 456. 457. 458. 459. 460. 461. 462. 463. 464. 465. 466. 467. 468. 469. 470. 471. 472. 473. 474. 475. 476. 477. 478. 479. 480. 481. 482. 483. 484. 485. 486. 487. 488. 489. 490. 491. 492. 493. 494. 495. 496. 497. 498. 499. 500. 501. 502. 503. 504. 505. 506. 507. 508. 509. 510. 511. 512. 513. 514. 515. 516. 517. 518. 519. 520. 521. 522. 523. 524. 525. 526. 527. 528. 529. 530. 531. 532. 533. 534. 535. 536. 537. 538. 539. 540. 541. 542. 543. 544. 545. 546. 547. 548. 549. 550. 551. 552. 553. 554. 555. 556. 557. 558. 559. 560. 561. 562. 563. 564. 565. 566. 567. 568. 569. 570. 571. 572. 573. 574. 575. 576. 577. 578. 579. 580. 581. 582. 583. 584. 585. 586. 587. 588. 589. 590. 591. 592. 593. 594. 595. 596. 597. 598. 599. 600. 601. 602. 603. 604. 605. 606. 607. 608. 609. 610. 611. 612. 613. 614. 615. 616. 617. 618. 619. 620. 621. 622. 623. 624. 625. 626. 627. 628. 629. 630. 631. 632. 633. 634. 635. 636. 637. 638. 639. 640. 641. 642. 643. 644. 645. 646. 647. 648. 649. 650. 651. 652. 653. 654. 655. 656. 657. 658. 659. 660. 661. 662. 663. 664. 665. 666. 667. 668. 669. 670. 671. 672. 673. 674. 675. 676. 677. 678. 679. 680. 681. 682. 683. 684. 685. 686. 687. 688. 689. 690. 691. 692. 693. 694. 695. 696. 697. 698. 699. 700. 701. 702. 703. 704. 705. 706. 707. 708. 709. 710. 711. 712. 713. 714. 715. 716. 717. 718. 719. 720. 721. 722. 723. 724. 725. 726. 727. 728. 729. 730. 731. 732. 733. 734. 735. 736. 737. 738. 739. 740. 741. 742. 743. 744. 745. 746. 747. 748. 749. 750. 751. 752. 753. 754. 755. 756. 757. 758. 759. 760. 761. 762. 763. 764. 765. 766. 767. 768. 769. 770. 771. 772. 773. 774. 775. 776. 777. 778. 779. 780. 781. 782. 783. 784. 785. 786. 787. 788. 789. 790. 791. 792. 793. 794. 795. 796. 797. 798. 799. 800. 801. 802. 803. 804. 805. 806. 807. 808. 809. 810. 811. 812. 813. 814. 815. 816. 817. 818. 819. 820. 821. 822. 823. 824. 825. 826. 827. 828. 829. 830. 831. 832. 833. 834. 835. 836. 837. 838. 839. 840. 841. 842. 843. 844. 845. 846. 847

1. SPONSORING MILITARY ACTIVITY

10. Διοικητική

DD FORM 1473 (PAGE 1)

DD FORM 1473

S. A. 0101. 307. 0001

Security Classification

Unclassified

Security Classification

| 14 KEY WORDS | LINK A | | LINK B | | LINK C | |
|--|--------|----|--------|----|--------|----|
| | ROLE | WT | ROLE | WT | ROLE | WT |
| Game theory Cooperative games Kernel Bargaining sets Nucleolus Transfer schemes Global stability Asymptotic stability Characteristic function games Dynamical systems Differential equations | | | | | | |

Unclassified

Security Classification